



# Haloalkanes

1(a). This question is about the alcohols **A–F** shown below.



Alcohol **A** can be prepared by the alkaline hydrolysis of the bromoalkane,  $(\text{CH}_3)_2\text{CHCH}_2\text{CH}_2\text{Br}$ . The hydrolysis with aqueous  $\text{NaOH}$  is shown in **equation 5.1**.




A student gently heats a mixture of  $(\text{CH}_3)_2\text{CHCH}_2\text{CH}_2\text{Br}$  and  $\text{NaOH}(\text{aq})$  for 25 minutes.

- i. Calculate the atom economy for the preparation of alcohol **A** in **equation 5.1**.

atom economy = ..... % [2]

- ii. Outline the mechanism for the alkaline hydrolysis of  $(\text{CH}_3)_2\text{CHCH}_2\text{CH}_2\text{Br}$ . The structure of  $(\text{CH}_3)_2\text{CHCH}_2\text{CH}_2\text{Br}$  has been provided.

Show curly arrows, relevant lone pairs and dipoles, and the products.



[3]

- iii. Name this type of mechanism.

[1]

## 4.2.2 Haloalkanes

- (b). The student decides to prepare alcohol **A** using the same method as in the part above but using the chloroalkane  $(\text{CH}_3)_2\text{CHCH}_2\text{CH}_2\text{Cl}$  instead of the bromoalkane,  $(\text{CH}_3)_2\text{CHCH}_2\text{CH}_2\text{Br}$ .

State and explain how the rates of hydrolysis of the chloroalkane and the bromoalkane would differ.

-----  
-----  
-----  
-----  
-----


[2]

2. This question is about 1-iodopentane,  $\text{CH}_3\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{I}$ .

1-iodopentane can be hydrolysed by aqueous sodium hydroxide.

- i. Outline the mechanism for this reaction.

Include curly arrows, relevant dipoles and the final product(s).



[3]

- ii. 1-iodopentane can also be hydrolysed by water using aqueous silver nitrate, with ethanol as the solvent.

A student uses this method to compare the rates of hydrolysis of 1-iodopentane and 1-bromopentane.

What measurement and observation would allow the student to compare the rates of hydrolysis?

-----  
-----

[1]

- iii. 1-iodopentane was found to react faster than 1-bromopentane.

Explain why.

-----  
-----  
-----

[2]


3. An alcohol can be prepared by hydrolysing the haloalkane  $C_2H_5CHBrCH_3$  with aqueous sodium hydroxide.

i. Outline the mechanism for this reaction.

Show curly arrows and relevant dipoles.

[3]

- ii. The infrared (IR) spectrum for  $C_2H_5CHBrCH_3$  is shown in **Fig. 25.2**. The C–Br bond absorption is labelled.



**Fig. 25.2**

Outline how IR spectroscopy could be used to show that the bromoalkane functional group has reacted and that the alcohol functional group has formed.

---

---

---

---

[2]

## 4.2.2 Haloalkanes

**4(a).** This question is about the hydrolysis of haloalkanes.

The rate of hydrolysis of a haloalkane depends on the halogen present.

State and explain how the halogen in the haloalkane affects the rate of hydrolysis.


-----  
-----  
-----  
-----  
-----

[2]

**(b).** Chlorocyclohexane is hydrolysed with aqueous sodium hydroxide.

Outline the mechanism for this reaction.

Show curly arrows, relevant dipoles and the products.



[3]

**(c).** A student hydrolyses a haloalkane, **E**, using the following method.

- 0.0100 mol of haloalkane **E** is refluxed with excess NaOH(aq) to form a reaction mixture containing an organic product **F**.
- The reaction mixture is neutralised with dilute nitric acid.
- Excess AgNO<sub>3</sub>(aq) is added to the reaction mixture. 1.88 g of a precipitate **G** forms.

Organic product, **F**, has a molar mass of 74.0 g mol<sup>-1</sup> and has a chiral carbon atom.

- i. Draw a **labelled** diagram to show how the student would carry out the hydrolysis of haloalkane **E**.

[2]

## 4.2.2 Haloalkanes


- ii. Analyse the information to identify **E**, **F** and **G**.

Show your working.

[3]

5. Alcohols are used in organic synthesis.

Pentan-2-ol can be prepared by the alkaline hydrolysis of 2-iodopentane.



The reaction mixture is boiled for 20 minutes.

- i. State the most appropriate technique that could be used to boil the reaction mixture for 20 minutes.

----- [1]

- ii. Describe the mechanism for the alkaline hydrolysis of 2-iodopentane.


In your answer, include the name of the mechanism, curly arrows and relevant dipoles.

name of mechanism: .....

[4]

6. Organic compounds can be prepared in the laboratory using synthetic routes with two or more stages.

A student devises a two-stage synthesis of cyclohexene from bromocyclohexane.



## 4.2.2 Haloalkanes

- i. Suggest the structure of **intermediate E** and the reagent(s) and conditions for **step 2**.

reagent(s) and  
conditions

-----

- ii. The student carries out this synthesis and obtains 1.23 g of pure cyclohexene from 5.50 g of bromocyclohexane.

Calculate the percentage yield of cyclohexene.

Give your final answer to an **appropriate** number of significant figures.

percentage yield = ..... % **[3]**

## 4.2.2 Haloalkanes

**7(a).** Haloalkanes are hydrolysed by aqueous sodium hydroxide.

- i. Outline the mechanism of the reaction of 1-bromobutane with aqueous sodium hydroxide.

Include curly arrows, relevant dipoles and the structure of the organic product.

[3]

- ii. Name the type of mechanism in (i).

[1]

- iii. The organic product in (i) can be formed faster using a different haloalkane than 1-bromobutane.

Identify this haloalkane.

Explain your answer.

Haloalkane

.....

Explanation

.....

[1]

**(b).** The use of some haloalkanes, such as chlorotrifluoromethane, has been banned as they form  $Cl\cdot$  radicals which break down ozone.

- i. Construct an equation to show the formation of  $Cl\cdot$  radicals from chlorotrifluoromethane.

[1]

- ii. Ozone is broken down by  $Cl\cdot$  radicals in a two-step process.

Write the equations for the two steps and the overall equation for this process.

Step 1

.....

Step 2

.....

Overall equation

..... [3]







## 4.2.2 Haloalkanes

-----  
-----  
-----  
-----  
-----  
-----

[6]

- ii. Compound **G** is heated with compound **F** in the presence of a small amount of concentrated sulfuric acid to form organic compound **H**.

Draw the structure of the organic compound **H**.

[2]

9. In the stratosphere, nitrogen oxides can catalyse the breakdown of ozone.

- i. State **two** sources of nitrogen oxides in the stratosphere.

-----  
-----

[1]

- ii. Write equations to show how nitrogen monoxide catalyses the breakdown of ozone.

-----  
-----  
-----

[2]

10. Alcohols can be prepared from halogenoalkanes. 2,2-dimethylpropan-1-ol can be prepared by hydrolysis of a chloroalkane with aqueous sodium hydroxide.

i. Write the equation for this reaction.

Use structures for the organic compounds.


[1]

ii. Outline the mechanism for this reaction.

Show curly arrows and relevant dipoles.


[2]

11. Compound **B**, shown below, can be used to synthesise organic compounds with different functional groups.



**Compound B**

The structure of compound **F** is shown below.



**Compound F**

i. What is the empirical formula of compound **F**?


----- [2]

## 4.2.2 Haloalkanes

- ii. A student plans a two-stage synthesis for preparing compound **F** from compound **B**.

The synthesis first prepares compound **E**, as shown in the flowchart.

Draw the structure of compound **E** in the box and state the reagents for each stage on the dotted lines.



[3]

12. Nitrogen forms several different oxides.

$\text{N}_2\text{O}$  is a useful anaesthetic and  $\text{NO}$  has been linked to the depletion of ozone in the stratosphere.

$\text{NO}$  radicals catalyse the breakdown of ozone in the stratosphere.

Write **two** equations to show how  $\text{NO}$  radicals catalyse this breakdown.

.....  
..... [2]

- 13(a). Reaction mechanisms use curly arrows and can involve electrophiles and nucleophiles.

- i. What does a *curly arrow* represent in mechanisms?

.....  
..... [1]

- ii. What is meant by the term *nucleophile*?

.....  
..... [1]









[5]

- iii. Radical substitution produces a mixture of organic products.

Suggest **two** reasons why.

[2]

14. This question is about the compounds shown below.

**B****C****D****E****F****G****H**

Compound **G** was once used as a propellant in aerosols. Compound **G** has been linked with depletion of the ozone layer in the stratosphere.

- i. State **two** properties that made compound **G** suitable for use as an aerosol.

1


2

[1]



16(a). The flowchart shows how 2-methylbut-2-ene can be converted into a number of organic products.

Complete the flowchart by drawing an organic structure in the box below.



[1]

(b). Identify reagent **A**.

----- [1]

(c). In the flowchart, **reaction R** forms a mixture of two alcohols that are structural isomers of  $C_5H_{12}O$ .

i. State the reagents and conditions needed for **reaction R**.

----- [1]

ii. What is meant by the term *structural isomers*?

----- [1]

iii. Draw the two structural isomers of  $C_5H_{12}O$  formed in **reaction R**.

[2]

iv. Suggest why 2-methylbut-2-ene is less soluble in water than either of the structural isomers formed.

----- [2]



## 4.2.2 Haloalkanes

- 17(a).** 1-Bromobutane,  $\text{CH}_3\text{CH}_2\text{CH}_2\text{CH}_2\text{Br}$ , reacts with methoxide ions,  $\text{CH}_3\text{O}^-$ , by nucleophilic substitution.

Suggest how the methoxide ion can act as a nucleophile.

-----  
-----  
----- [1]

- (b).** Using the 'curly arrow' model, suggest the mechanism for this reaction.

Show any relevant dipoles.

[3]

- (c).** 1-Iodobutane also reacts with methoxide ions.

Indicate, by placing a tick in one of the boxes, how the use of 1-iodobutane would affect the rate of reaction compared with that of 1-bromobutane.

1-Iodobutane does not change the rate	
1-Iodobutane increases the rate	
1-Iodobutane decreases the rate	

Explain your answer.

-----  
-----  
----- [1]

- (d).** The ethanoate ion,  $\text{CH}_3\text{COO}^-$  acts as a nucleophile when reacting with 1-bromobutane in a substitution reaction.


Draw the skeletal formula and give the name of the organic product formed in this reaction.

skeletal formula

name of product ..... [2]

**18(a).** Ethers are a homologous series of organic compounds containing the R–O–R functional group.

The structures and names of two ethers are shown in **Fig. 4.1**.



**Fig. 4.1**

Draw the **skeletal** formula of the ether, 2-ethoxy-3-methylbutane.

[1]

**(b).** Ethers can be prepared by nucleophilic substitution of haloalkanes with alkoxide ions, RO<sup>−</sup>.

i. Alkoxide ions can be prepared by reacting sodium with an alcohol. A gas is also formed.

Write an equation for the formation of methoxide ions from sodium and an alcohol.

----- [1]

ii. Methoxyethane, shown in **Fig. 4.1**, can be prepared by reacting bromoethane, CH<sub>3</sub>CH<sub>2</sub>Br, with methoxide ions, CH<sub>3</sub>O<sup>−</sup>.

Suggest the mechanism for the nucleophilic substitution of CH<sub>3</sub>CH<sub>2</sub>Br with CH<sub>3</sub>O<sup>−</sup>.

Show curly arrows, charges, relevant dipoles, and products.

[3]

iii. In this mechanism, explain how CH<sub>3</sub>O<sup>−</sup> ions have acted as a nucleophile.

State the type of bond fission that takes place.

----- [1]

## 4.2.2 Haloalkanes

- (c). 2-Ethoxypropane, shown in **Fig. 4.1**, is analysed by  $^1\text{H}$  NMR spectroscopy.

Complete the table to predict the  $^1\text{H}$  NMR spectrum of 2-ethoxypropane.  
You may **not** need to use all the rows.

Chemical shift, $\delta/\text{ppm}$	Relative peak area	Splitting pattern

[4]

- (d). In organic reactions, alkoxide ions can also act as a base.

The diagram below shows an incomplete mechanism for the reaction of a diester with methoxide ions,  $\text{CH}_3\text{O}^-$  (**Step 1**), followed by reaction of the intermediate with bromoethane (**Step 2**).

- i. For **Step 1**, add curly arrows to show how  $\text{CH}_3\text{O}^-$  reacts with the diester to form the intermediate.  
In the box, draw the structure of the organic product formed in **Step 2**.



[3]

- ii. Explain how  $\text{CH}_3\text{O}^-$  ions have acted as a base in this mechanism.

-----

----- [1]

## 4.2.2 Haloalkanes

19. Haloalkanes can undergo hydrolysis.

A student carries out an experiment to find the relative rate of hydrolysis of 1-chloropropane,  $C_3H_7Cl$ , 1-bromopropane,  $C_3H_7Br$ , and 1-iodopropane,  $C_3H_7I$ .

The student adds  $2\text{ cm}^3$  of ethanol to  $2\text{ cm}^3$  of aqueous silver nitrate to three test tubes labelled **A**, **B** and **C**.

The student adds 5 drops of a different haloalkane to each test-tube in rapid succession and shakes each tube. The student measures the time for a precipitate to form in each test-tube.

The results are shown below.

Test tube	Haloalkane	Time taken for reaction to take place
<b>A</b>	$C_3H_7Cl$	about half an hour
<b>B</b>	$C_3H_7Br$	a few minutes
<b>C</b>	$C_3H_7I$	a few seconds

- i. Write an **ionic** equation involving aqueous silver nitrate for formation of **one** of the precipitates.

-----  
[1]

- ii. What do the experimental results tell you about the carbon–halogen bond enthalpies?

-----  
[1]

- iii. How could the student modify their experiment so that it could be completed in less time?

-----  
[1]

**END OF QUESTION PAPER**