


How Fast

1. A student investigates the reaction between ethanoic acid, $\text{CH}_3\text{COOH}(\text{l})$ and methanol, $\text{CH}_3\text{OH}(\text{l})$, in the presence of an acid catalyst. The equation is shown below.




The student carries out an experiment to determine the order of reaction with respect to CH_3COOH .

The student uses a large excess of CH_3OH . The temperature is kept constant throughout the experiment.

The student takes a sample from the mixture every 20 minutes, and then determines the concentration of the ethanoic acid in each sample.

From the experimental results, the student plots the graph below.



- i. Explain why the student uses a large excess of methanol in this experiment.

[1]

- ii. Use the half-life of this reaction to show that the reaction is first order with respect to CH_3COOH .

Show your working on the graph and below.


[2]

- iii. Determine the initial rate of reaction.

initial rate = $\text{mol dm}^{-3} \text{ min}^{-1}$ [2]

5.1.1 How Fast


2. Three students carry out a rates investigation on the reaction between bromine and propanone in the presence of hydrochloric acid.




Each student investigates the effect of changing the concentration of one of the reactants whilst keeping the other concentrations constant.

Their results are shown below.

Results of student 1



Results of student 2




Results of student 3

Experiment	$[\text{Br}_2(\text{aq})] / \text{mol dm}^{-3}$	$[\text{CH}_3\text{COCH}_3(\text{aq})] / \text{mol dm}^{-3}$	$[\text{H}^+(\text{aq})] / \text{mol dm}^{-3}$	Initial rate / $10^{-5} \text{ mol dm}^{-3} \text{ s}^{-1}$
1	0.004	1.60	0.20	1.25
2	0.004	1.60	0.40	2.50

3. This question is about organic reactions.

Compound **A** is formed when ethanal is mixed with OH^- (aq) ions, which act as a catalyst.

The balanced equation is shown in **reaction 6.1** below.



- i. Give the systematic name for compound **A**.

----- [1]


- ii. What type of reaction has taken place?

----- [1]

- iii. **Reaction 6.1** takes place in two steps. OH^- ions act as a catalyst.

In **step 1**, ethanal reacts with OH^- ions to set up an acid–base equilibrium.
In **step 2**, compound **A** is formed.

- Complete the equilibrium for **step 1** and label the conjugate acid–base pairs as: **A1**, **B1** and **A2**, **B2**.



- Suggest the equation for **step 2**.

[3]

- iv. A similar reaction takes place when propanone, $(\text{CH}_3)_2\text{CO}$, is mixed with OH^- (aq) ions.


Draw the structure of the organic product of this reaction.

[1]

5.1.1 How Fast

- 4(a).** A student carries out an investigation to find the activation energy, E_a , and the pre-exponential factor, A , of a reaction.

The student determines the rate constant, k , at different temperatures, T .
The student then plots a graph of $\ln k$ against $1/T$ as shown below.



- i. Draw a best-fit straight line and calculate the activation energy, in J mol^{-1} .
Give your answer to **three** significant figures.

Show your working.

activation energy, $E_a = + \dots\dots\dots \text{J mol}^{-1}$ **[3]**


- ii. Use the graph to calculate the value of the pre-exponential factor, A .

Show your working.

pre-exponential factor, $A = \dots\dots\dots$ **[2]**

5.1.1 How Fast

5(a). Aqueous solutions of hydrogen peroxide, $\text{H}_2\text{O}_2(\text{aq})$, decompose as in the equation below.



A student investigates the decomposition of $\text{H}_2\text{O}_2(\text{aq})$ by measuring the volume of oxygen gas produced over time. All gas volumes are measured at room temperature and pressure.


The student uses 25.0 cm^3 of $2.30 \text{ mol dm}^{-3} \text{ H}_2\text{O}_2$.

From the results, the student determines the concentration of $\text{H}_2\text{O}_2(\text{aq})$ at each time. The student then plots a concentration–time graph.

Suggest a different experimental method that would allow the rate of this reaction to be followed over time.

[1]

(b). Aqueous solutions of hydrogen peroxide, $\text{H}_2\text{O}_2(\text{aq})$, decompose as in the equation below.



A student investigates the decomposition of $\text{H}_2\text{O}_2(\text{aq})$ by measuring the volume of oxygen gas produced over time. All gas volumes are measured at room temperature and pressure.

The student uses 25.0 cm^3 of $2.30 \text{ mol dm}^{-3} \text{ H}_2\text{O}_2$.

From the results, the student determines the concentration of $\text{H}_2\text{O}_2(\text{aq})$ at each time. The student then plots a concentration–time graph.

5.1.1 How Fast


- (c). Determine the total volume of oxygen, measured at room temperature and pressure, that the student should be prepared to collect in this investigation.

Suggest apparatus that would allow this gas volume to be collected, indicating clearly the scale of working.

[3]

6. This question is about reactions of hydrogen peroxide, H₂O₂.

Hydrogen peroxide, H₂O₂, iodide ions, I⁻, and acid, H⁺, react as shown in the equation below.



A student carries out several experiments at the same temperature, using the initial rates method, to determine the rate constant, *k*, for this reaction.

The results are shown below.

Experiment	Initial concentrations			Rate / 10 ⁻⁶ mol dm ⁻³ s ⁻¹
	[H ₂ O ₂ (aq)] / mol dm ⁻³	[I ⁻ (aq)] / mol dm ⁻³	[H ⁺ (aq)] / mol dm ⁻³	
1	0.0100	0.0100	0.100	2.00
2	0.0100	0.0200	0.100	4.00
3	0.0200	0.0100	0.100	4.00
4	0.0200	0.0100	0.200	4.00

- i. Determine the rate equation and calculate the rate constant, *k*, including units.

k = units [3]


5.1.1 How Fast

- ii. The rate constant, k , for this reaction is determined at different temperatures, T .

Explain how the student could determine the activation energy, E_a , for the reaction graphically using values of k and T .

[3]

- 7(a). A student investigates the rate of reaction between iodine, I_2 , and propanone, CH_3COCH_3 , in the presence of H^+ ions. The student uses $HCl(aq)$ to supply H^+ ions.



The student follows the method outlined below.

1. The student starts the reaction by mixing the following solutions.

1.00 cm³ of 1.00 mol dm⁻³ $I_2(aq)$
49.5 cm³ of 1.00 mol dm⁻³ $CH_3COCH_3(aq)$
49.5 cm³ of 1.00 mol dm⁻³ $HCl(aq)$

2. The student places a sample of the reaction mixture in a colorimeter, immediately starts a stopwatch, and records the absorbance.

3. The student records the absorbance every 100 s. The results are shown below.

Time/s	Absorbance
0	0.80
100	0.67
200	0.51
300	0.44
400	0.28
500	0.18
600	0.05

Explain why absorbance decreases during the experiment.

[1]

5.1.1 How Fast


- (b). Absorbance is proportional to the concentration of I_2 .

Calculate the concentration of I_2 at the start of the experiment and after 500 s.

Time/s	Absorbance	$[I_2(aq)]/\text{mol dm}^{-3}$
0	0.80	
500	0.18	

[2]

- (c). i. Plot a graph of absorbance against time and draw a line of best fit.



[3]

- ii. Use your graph to find the order of reaction with respect to iodine.
Explain your reasoning.

5.1.1 How Fast

Order


.....

Explanation

.....

----- **[2]**

(d). A three step mechanism has been proposed for this reaction.




Complete the mechanism by adding equations for **Step 1** and **Step 3** in the boxes below.

Step 1 (slow)	
Step 2 (fast)	$ \begin{array}{c} \text{OH} \\ \\ \text{H}_3\text{C}-\text{C}^+-\text{CH}_3 \\ \\ \text{O} \end{array} \longrightarrow \begin{array}{c} \text{OH} \\ \\ \text{H}_3\text{C}-\text{C}=\text{CH}_2 \end{array} + \text{H}^+ $
Step 3 (fast)	

[2]

8. This question is about redox reactions.

*Bromine, Br₂, is formed in the redox reaction shown below.



A student plans an investigation, using the initial rates method, to determine the rate equation and rate constant for this reaction.

The student is supplied with solutions containing the following:

- 0.300 mol dm⁻³ Br⁻(aq)
- 0.300 mol dm⁻³ BrO₃⁻(aq)
- 0.300 mol dm⁻³ H⁺(aq).

The student is also supplied with distilled water and normal laboratory glassware.

The student uses a total volume of 30 cm³ for each experiment and measures the initial rate of formation of Br₂(aq).


The results of the student's experiments are shown below.

5.1.1 How Fast

Experiment	$[\text{Br}^-(\text{aq})] / \text{mol dm}^{-3}$	$[\text{BrO}_3^-(\text{aq})] / \text{mol dm}^{-3}$	$[\text{H}^+(\text{aq})] / \text{mol dm}^{-3}$	Initial rate / $10^{-3} \text{ mol dm}^{-3} \text{ s}^{-1}$
1	0.100	0.100	0.100	1.20
2	0.025	0.100	0.100	0.30
3	0.100	0.050	0.100	0.60
4	0.100	0.050	0.050	0.15

Show how the student could obtain the concentrations for experiments 1–4 and determine the rate constant for this reaction.

9(a). Hydrogen peroxide reacts with iodide ions in acid conditions, as shown below.



A student investigates the rate of this reaction by carrying out four experiments at the same temperature. The student's results are shown below.

Experiment	[H ₂ O ₂ (aq)] / mol dm ⁻³	[I ⁻ (aq)] / mol dm ⁻³	[H ⁺ (aq)] / mol dm ⁻³	Initial rate / mol dm ⁻³ s ⁻¹
1	0.0010	0.20	0.10	5.70 × 10 ⁻⁶
2	0.0020	0.20	0.10	1.14 × 10 ⁻⁵
3	0.0020	0.20	0.20	1.14 × 10 ⁻⁵
4	0.0040	0.40	0.10	4.56 × 10 ⁻⁵

The rate equation is: $rate = k [\text{H}_2\text{O}_2(\text{aq})] [\text{I}^-(\text{aq})]$

- Show that the student's results support this rate equation.
- Calculate the rate constant, *k*, for this reaction.

Give your answer to **two** significant figures, in standard form and with units.



In your answer you should make clear how the experimental results provide evidence for the rate equation.

(b). The student concluded that $\text{H}^+(\text{aq})$ ions act as a catalyst.

Explain why the student's conclusion is **not** correct.

[1]

(c). A four-step mechanism has been proposed for this reaction.
The rate-determining step is the first step.

i. State what is meant by the term *rate-determining step*.

[1]

ii. The equation for **Step 3** in the four-step mechanism is shown below.

Suggest equations for the other three steps.
State symbols are **not** required.

Step 1:

Step 2:


Step 3: $\text{HIO} + \text{I}^- \rightarrow \text{I}_2 + \text{OH}^-$

Step 4:

[3]

5.1.1 How Fast


10(a). Hydrogen, H_2 , reacts with nitrogen monoxide, NO , as shown below:



The rate equation for this reaction is:

$$\text{rate} = k[\text{H}_2(\text{g})][\text{NO}(\text{g})]^2$$

The concentration of $\text{NO}(\text{g})$ is changed and a rate–concentration graph is plotted.



The chemist uses $\text{H}_2(\text{g})$ of concentration $2.0 \times 10^{-2} \text{ mol dm}^{-3}$.

Using values from the graph, calculate the rate constant, k , for this reaction.

Give your answer to **two** significant figures and in **standard form**.

Show your working.

$k = \dots\dots\dots$ units $\dots\dots\dots$ [4]

(b). A chemist investigates the effect of changing the concentration of $\text{H}_2(\text{g})$ on the initial reaction rate at two different temperatures.

The reaction is first order with respect to $\text{H}_2(\text{g})$.


i. Using the axes below, sketch **two** graphs of the results.

Label the graphs as follows:

○ L for the lower temperature

5.1.1 How Fast

- **H** for the higher temperature.




[2]

- ii. State the effect of the higher temperature on the rate constant, k .

----- [1]

- (c). The reaction can also be shown as being first order with respect to $\text{H}_2(\text{g})$ by continuous monitoring of $[\text{H}_2(\text{g})]$ during the course of the reaction.


- Using the axes below, sketch a graph to show the results.
- State how you would use the graph to show this first order relationship for $\text{H}_2(\text{g})$.



----- [2]

5.1.1 How Fast

(d). The chemist proposes a three-step mechanism for the reaction:



i. On the dotted line below, write the equation for step 3.


step 1:	$2\text{NO} \rightarrow \text{N}_2\text{O}_2$	fast
step 2:	$\text{H}_2 + \text{N}_2\text{O}_2 \rightarrow \text{N}_2\text{O} + \text{H}_2\text{O}$	slow
step 3:	fast

[1]

ii. Explain why this mechanism is consistent with the rate equation $\text{rate} = k[\text{H}_2(\text{g})][\text{NO}(\text{g})]^2$.

[1]

11(a). A student carries out an initial rates investigation on the reaction below.



From the results, the student determines the rate equation for this reaction:

$$\text{rate} = k [\text{I}^-(\text{aq})]^2 [\text{IO}_3^-(\text{aq})] [\text{H}^+(\text{aq})]^2$$

i. What is the overall order of reaction?

----- [1]


ii. A proposed mechanism for this reaction takes place in several steps.

Suggest **two** reasons why it is unlikely that this reaction could take place in one step.

[2]

5.1.1 How Fast

- (b). On the rate—concentration graphs below, sketch lines to show the relationship between initial rate and concentration for IO_3^- (aq) and H^+ (aq).



[2]

- (c). The table below shows some of the student's results.

- i. Complete the table by adding the missing initial rates in the boxes.

	$[\text{I}^-](\text{aq}) / \text{mol dm}^{-3}$	$[\text{IO}_3^-](\text{aq}) / \text{mol dm}^{-3}$	$[\text{H}^+](\text{aq}) / \text{mol dm}^{-3}$	Initial rate / $\text{mol dm}^{-3} \text{ s}^{-1}$
Experiment 1	0.015	0.010	0.020	0.60
Experiment 2	0.045	0.010	0.020	
Experiment 3	0.060	0.040	0.080	

[2]

- ii. Calculate the rate constant, k , for this reaction. Include units.

Give your answer to **two** significant figures.

$$k = \dots\dots\dots \text{units} \dots\dots\dots [3]$$

- iii. The student repeats Experiment 1 using $0.020 \text{ mol dm}^{-3}$ methanoic acid, $\text{HCOOH}(\text{aq})$ ($pK_a = 3.75$), instead of $0.020 \text{ mol dm}^{-3} \text{ HCl}(\text{aq})$ as a source of $\text{H}^+(\text{aq})$.

Determine the initial rate in this experiment. Show your working.

$$\text{initial rate} = \dots\dots\dots \text{mol dm}^{-3} \text{ s}^{-1} [3]$$

12. This question is about numbers and patterns in chemistry.

This question looks at number relationships. For calculations, show your working.

- i. What is the oxidation number of nitrogen in each species?

N_2O_4 NO_3^- NH_4^+

- ii. [1]
 iii. What mass of KMnO_4 is needed to prepare a 250.0 cm^3 solution with a concentration of $0.200 \text{ mol dm}^{-3} \text{ KMnO}_4$?


mass = g [2]

- iv. What are the units of the rate constant for a reaction with an overall order of 3?

units = [1]

- v. How many molecules are in 38.25 g of tartaric acid?

Give your answer to an **appropriate** number of significant figures and in standard form.




tartaric acid

number of molecules = [2]

5.1.1 How Fast


13. Methanoic acid and bromine react as in the equation below.



A student investigates the rate of this reaction by monitoring the concentration of bromine over time.

The student uses a large excess of HCOOH to ensure that the order with respect to HCOOH will be effectively zero.

From the experimental results, the student plots the graph below.



- a. Suggest how the concentration of the bromine could have been monitored.

----- [1]

- b. Suggest a different experimental method that would allow the rate of this reaction to be followed over time.

----- [1]

5.1.1 How Fast

- c. Why would use of excess HCOOH ensure that the order with respect to HCOOH is effectively zero?

[1]

- d. * Using the graph, determine
- the initial rate of reaction
 - the rate constant.

Your answer must show full working using the graph and the lines below as appropriate.

[6]

5.1.1 How Fast

- (b). A student carries out an investigation to find the activation energy, E_a , of a reaction.

From the results, the student determines the rate constant, k , at different temperatures, T .

The student then calculates $1/T$ and $\ln k$, as shown in **Table 19.1**.


Temperature, T / K	Rate constant, k / s^{-1}	$1 / T / K^{-1}$	$\ln k$
278	1.50×10^{-6}	3.60×10^{-3}	-13.41
290	1.51×10^{-5}		-11.10
298	4.11×10^{-5}	3.34×10^{-3}	-10.10
308	1.99×10^{-4}	3.23×10^{-3}	
323	1.40×10^{-3}	3.10×10^{-3}	-6.57

Table 19.1

Add the missing values to **Table 19.1** and plot a graph of $\ln k$ against $1/T$ on the graph paper opposite.

Using your graph, calculate the activation energy of the reaction.


Show your working.



activation energy, $E_a = \dots\dots\dots$ kJ mol^{-1} **[4]**

5.1.1 How Fast

15. Iodine monochloride, ICl, can react with hydrogen to form iodine.



This reaction was carried out several times using different concentrations of ICl or H₂. The initial rate of each experiment was calculated and the results are shown below. Initial concentrations are shown for each experiment.

	[ICl] / mol dm ⁻³	[H ₂] / mol dm ⁻³	Rate / mol dm ⁻³ s ⁻¹
Experiment 1	0.250	0.500	2.04 × 10 ⁻²
Experiment 2	0.500	0.500	4.08 × 10 ⁻²
Experiment 3	0.125	0.250	5.10 × 10 ⁻³

- i. Calculate the rate constant, *k*, for this reaction. Include units in your answer.

Show **all** your working.

$$k = \dots\dots\dots \text{units} \dots\dots\dots \text{ [4]}$$

- ii. Calculate the rate of reaction when ICl has a concentration of 3.00 × 10⁻³ mol dm⁻³ and H₂ has a concentration of 2.00 × 10⁻³ mol dm⁻³.

Show **all** your working.

$$\text{rate} = \dots\dots\dots \text{ mol dm}^{-3} \text{ s}^{-1} \text{ [1]}$$

END OF QUESTION PAPER