
Amines

1. This question is about organic compounds containing nitrogen.

Salt H, $(CH_3)_2CHNH_3CI$, is used in the manufacture of garden weedkillers.

The flowchart shows the synthesis of the salt \mathbf{H} from propanone.

Complete the flowchart. Show structures for organic compounds.



[5]

2. This question is about organic compounds containing nitrogen.

Sodium cyanide, NaCN, can be reacted with many organic compounds to increase the length of a carbon chain.


i. 1-Chloropropane, CH₃CH₂CH₂C*I*, reacts with ethanolic sodium cyanide by nucleophilic substitution.

Outline the mechanism for this reaction.

Include curly arrows, relevant dipoles and the structure of the organic product.

ii. Compound **G** is used to synthesise compounds **H** and **I** as shown in the flowchart below.

Complete the flowchart showing the structure of compound **G** and the **formulae** of the reagents for **Reaction 2** and **Reaction 3**.



iii. Compound **H** reacts with dilute hydrochloric acid to form a salt.

Explain why compound ${\bf H}$ can react with dilute hydrochloric acid and suggest a structure for the salt formed.


Explanation			

Structure

iv.	Compound I is the monomer for the biodegradable polymer J. Draw two repeat units of polymer J and suggest a reason why it is biodegradable.					
	[3]					

[2]


- **3.** A student synthesises 3-bromophenylamine, shown below, starting from nitrobenzene.
 - i. Complete the flowchart showing the structure of the intermediate and the **formulae** of the reagents for each stage.



4. A student plans a two-stage synthesis of alanine from lactic acid, CH₃CH(OH)COOH.

The synthesis first prepared compound H, as shown in the flowchart.

Draw the structure of compound ${\bf H}$ in the box and add the formulae of the reagents for each stage on the dotted lines.



[3]

5(a). A chemistry teacher carries out an experiment to synthesise 2-aminopropan-1-ol, CH₃CH(NH₂)CH₂OH.

The teacher asks a university chemistry department to test the 2-aminopropan-1-ol using proton NMR spectroscopy and mass spectrometry.


i. For the ¹H NMR analysis, the sample was dissolved in D₂O.

Complete the table to predict the ¹H NMR spectrum of CH₃CH(NH₂)CH₂OH after dissolving in D_2O .

¹ H NMR spectrum for CH ₃ CH(NH ₂)CH ₂ OH, dissolved in D ₂ O						
Chemical shift, δ/ ppm	Relative peak area	Splitting pattern				

[3]

ii. The mass spectrum for CH₃CH(NH₂)CH₂OH is shown below.




Give the formulae for the species responsible for **peak 1** and **peak 2** in the mass spectrum.

peak 1

peak 2

[1]

(c). In a separate experiment, the chemistry teacher prepares compound E from 2-aminopropan-1-ol.




i. One of the functional groups in compound **E** is a phenol.

Name the other functional groups in compound E.


.....[1]

ii. Draw the structures of the **two** organic products formed when compound **E** is heated under reflux with dilute hydrochloric acid.



[2]

6(a). This question is about the reactions of compounds with more than one functional group.A chemist investigates some reactions of compound L, as shown in the flowchart below.Complete the flowchart by showing the missing organic structures in the boxes.




[3]

(b). Outline the mechanism that occurs in **Reaction 1**. Include curly arrows, relevant dipoles and the name of the mechanism.

name of mechanism[4]

(c). The chemist synthesises compound \mathbf{M} , which can undergo both addition and condensation polymerisation.






i. Draw the repeat unit of the addition polymer formed from compound M.

[1]

ii. Draw two repeat units of the condensation polymer formed from compound M.

7. 4-Nitrobenzoic acid is an important compound in chemical synthesis. The flowchart below shows a synthesis involving 4-nitrobenzoic acid.



i. State suitable reactant(s) and conditions for step 1.

[1]

ii. In **step 2**, the –NO₂ group in compound **F** is reduced by tin and concentrated hydrochloric acid.

Write an equation for the reduction of compound F.

Show the structures of any organic compounds involved.

[2]

END OF QUESTION PAPER